

**Kills the
Fleas While it
Kills the Rodent**

Kaput[®]

COMBO BAIT[®] PELLETS

This product may be used only inside and within 50 feet of buildings or inside transport vehicles (ships, trains and aircraft).



**PLACE
PACKS**

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE & VOLES

NET WT. 4 LBS. (32 PACKS x 2 oz.)

ACTIVE INGREDIENTS:

Imidacloprid (CAS Number 138261-41-3) ... 0.020%

Warfarin (CAS Number 81-81-2) 0.025%

OTHER INGREDIENTS 99.955%

TOTAL 100.000%

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

HARMFUL IF SWALLOWED.

**SEE BACK PANEL FOR FIRST AID AND ADDITIONAL
PRECAUTIONARY STATEMENTS.**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This label must be in the user's possession at the time of product application.

Apply this product only for the sites, pests and application methods specified on this label.

If sufficient quantities of this bait are ingested, the use of this product kills fleas that commonly infest listed rodent hosts, including *Xenopsylla cheopis*.

Read This Label:

Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in locations out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and dogs, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, as well as unused, spoiled and unconsumed bait, as specified on this label.

Tamper-resistant bait stations must be used if children, pets, non-target mammals or birds have access to bait.

RATS AND MICE

Use Restrictions: Use this product to kill only Norway rats, roof rats and house mice, and their infested fleas, in and around homes and other residential buildings, industrial/agricultural buildings and similar man-made structures, and inside transport vehicles. Do not apply farther than 50 feet from buildings. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into direct contact with food. Do not use in edible product areas of feed processing plants, restaurants or other areas where food or feed is commercially prepared or processed. Do not contaminate food/feed or food/feed-handling equipment or place near or inside ventilation duct openings. Do not use this product to control listed species

within the occupied habitats of threatened or endangered species. Wear gloves when handling bait or animal carcasses. Do not sell this product in individual containers holding less than 4 pounds of bait.

Selection of Treatment Areas: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors or walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Norway Rats and Roof Rats: Apply 2 to 8 place packs per bait placement. Space placements at intervals of 15 to 30 feet in infested area. Maintain an uninterrupted supply of fresh bait for at least 10 days or until there no longer are signs of new feeding by rats.

House Mice: Open pack and apply 1/4 to 1/2 ounce of bait per placement. Space bait placements at intervals of 8 to 12 feet in infested area. At points of extremely high mouse activity use one 2-ounce pack. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of feeding by house mice cease to appear.

Follow-up (Rats & Mice): Immediately replace contaminated or spoiled bait. While wearing gloves, properly collect and dispose of all dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations. Regularly check the stations and replace bait as needed.

VOLES

Use Restrictions: Use this product to kill only meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetus*) and montane voles (*M. montanus*) and their infested fleas around (usually within 15 feet of) homes, and industrial, commercial, agricultural and similar man-made structures. Do not apply farther than 50 feet from buildings. Do not use this product to control listed species within the occupied habitats of threatened or endangered species. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not sell this product in individual containers holding less than 4 pounds of bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce (half of a pack) of bait into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. Check

placements daily. Remove any fouled or spoiled bait and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are signs of vole activity and feeding on bait.

Follow-up: While wearing gloves, collect and properly dispose of any carcasses and leftover bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Non-refillable container; do not reuse or refill this container. Offer for recycling, if available or reconditioning, if appropriate; otherwise, dispose of empty container in a sanitary landfill or by incineration.

Lot Number: See Container Lid

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves (PPE).

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are given, use detergent and hot water. Keep and wash PPE separately from other laundry.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Poison Information Center at 1-800-858-7378 for emergency medical treatment information.

If swallowed, immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes,** hold eye open and rinse slowly and gently

with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Treatment for Pet Poisoning

If animal eats bait, call veterinarian at once.

Notice to Physician or Veterinarian

Contains the anticoagulant Warfarin. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. For humans or animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait.

Notice: This product contains a chemical known to the state of California to cause birth defects or other reproductive harm.

WARRANTY

To the extent consistent with applicable law, Manufacturer and Seller make no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S. Patent No. 7,943,160 B2

Manufactured by:

Scimetrics
LTD. CORP.

New Solutions to Old Problems

P.O. Box 1045 Wellington, CO 80549
(970) 482-1330

Label Version 5/25/12

